# Journal of Basic and Applied Scientific Research (JBASR)



## An International Peer-reviewed journal

Number of issues per year: 12 ISSN: 2090-4304 (Print) ISSN: 2090-424x (Online)

Volume 1, Issue 2, November 2022



## J. Basic Appl. Sci. Res., Vol.12 No. 2: pp. 1-14, Year 2022

<b>Journal of Basic</b> a Monthly Publicatio	and Applied Scientific Res n	search (JBASR)				
	Number of issues per year: 12 ISSN: 2090-4304 (Print) ISSN: 2090-424x (Online)					
Journal of Taxic and Applied Scientific Research	<b>Journal of Basic and Applied Scientific Research (JBASR)</b> is a peer reviewed, open access international scientific journal dedicated for rapid publication of high quality original research articles as well as review articles in the all areas of basic and applied sciences.					
Trollad Seals Defeaters	Journal of Basic and Appli original and significant researc	) is devoted to the rapid publication of				
	Accounts	Energy	Nuclear Engineering			
	Agricultural Sciences	Engineering, All Fields	Oceanography			
Coopo	Applied Biology	Entomology	Oncology			
Scope	Biochemistry	Environment	Parasitology			
	Biological Sciences	Evolution	Petroleum & Gas			
	Biophysics	Fisheries	Pharmacology			
	Business and Economics	Food & Food Technology	Physics			
	Cell Biology	Genetics	Physiology			
	Chemical Engineering	Genomics	Plant Biology			
	Chemical Engineering	Geology	Population Biology			
	Chemistry	Immunology	Religious Studies			
	Civil Engineering	Infectious Diseases	Robotics			
	Civil Engineering	Law	Signal Transduction			
	Commerce	Marine Sciences	Social Sciences			
	Communication & IT	Marine Technology	Solid State Technology			
	Computer Science	Mathematics & Statistics	Space Science			
	Construction	Medical Technology	Textile Industry & Fabrics			
	Dentistry	Medicine	Toxicology			
	Developmental Biology	Microbiology	Transportation			
	Ecology	Nanotechnology	Veterinary Science			
	Endocrinology	Neuroscience	Zoology			

## **Editorial Board**

#### **Editor -in-Chief**

#### William Ebomoyi

Ph.D., Professor, Department of Health Studies, College of Health Sciences, Chicago State University, USA. E-mail: editor@textroad.com

#### **Associate Editors**

#### Prof. Dr. Sarwoko Mangkoedihardjo

Professor, Professional Engineer of Indonesian Society of Sanitary and Environmental Engineers, Indonesia

#### Saeid Chekani Azar

PhD of Veterinary Physiology; Faculty of Veterinary, Department of Physiology, Ataturk University, Erzurum 25010, **Turkey**.

#### Prof. Dr. Ashraf Latif Tadross

Head of Astronomy Department, Professor of Star Clusters and Galactic Structure, National Research Institute of Astronomy & Geophysics (NRIAG), 11421 Helwan, Cairo, **Egypt**.

#### Prof. Dr. Mario Bernardo-Filho

Full Professor, Universidade do Estado do Rio de Janeiro, Head, Laboratório de Radiofarmácia Experimental, Brazil.

#### Dr. Mounir M. Salem-Bekhet

Associate Professor of Microbiology, Department of Pharmaceutics, King Saud University, KSA.

#### Dr. Sandra Pacios Pujado

University of Pennsylvania, Philadelphia, PA, USA.

#### Vishal Patil, PhD

Materials Research Laboratory, University of California, Santa Barbara, CA, USA.

#### Dr. YUBAO CUI

Associate Professor, Department of Laboratory Medicine, Yancheng Health Vocational & Technical College, Jiangsu Province, P. R. China

#### Raja S Payyavula

Research Associate, Bio Science Division, Oak Ridge National Laboratory, Oak Ridge, Tennessee, USA.

#### **Dr. Zhihong Song**

The Ames Laboratory of US DOE, 2238 MBB Iowa State University, IA 54411 USA.

#### Prof. Dr. Nasser Fegh-hi Farahmand

Associate professor, Department of Industrial Management, Tabriz Branch, Islamic Azad University, Tabriz, Iran

#### Prof. Dr. Valdenir José Belinelo

Department of Health Sciences and Postgraduate Program in Tropical Agriculture, Federal University of Espirito Santo (UFES, São Mateus, ES, **Brazil** 

#### Dr. Chandrasekar Raman

Research Associate, Department of Biochemistry & Molecular Biophysics, Biotechnology Core Facility, 238, Burt Hall, Kansas State University, Manhattan 66506, KS, USA.

#### Mr. Jiban Shrestha

Scientist (Plant Breeding and Genetics), Nepal Agricultural Research Council, National Maize Research Program, Rampur, Chitwan, Nepal

#### Dr. Nadeem Javaid

Ph.D. (University of Paris-Est, France), Assistant Professor, Center for Advanced Studies in Telecommunications (CAST), COMSATS Institute of IT, Islamabad, **Pakistan** 

#### Dr. Syamkumar Siv Pillai

Program Manager-National Clean Plant Network - Fruit Trees, Washington State University, USA

#### Dr. Hala Ahmed Hafez Kandil

Professor Assistant, National Research Centre, Plant Nutrition Department. Dokki, Giza, Cairo, Egypt

#### Prof. Dr. Aziza Sharaby

Pests and Plant Protection Department, National Research Center, Cairo, Egypt

#### Prof. Dr. Sanaa T. El-Sayed

Ex Head of Biochemistry Department, Professor of Biochemistry, Genetic Engineering &Biotechnology Division, National Research Centre, **Egypt** 

#### Dr. Pratap V. Naikwade

M.Sc., Ph.D. Head, Department. of Botany, ASP College, Devrukh. Maharashtra, India.

#### Shadia M. Abdel-Aziz

Microbial Chemistry, National Research Center, Egypt

#### Dr. Tarig Osman Khider

Associate Professor, University of Bahri-Sudan, College of Applied and Industrial Sciences, Department of Pulp and Paper Technology, **Sudan** 

#### Dr. Hayman Z. Metwally

Associate Professor of Space Science cairo University **Egypt** and Vice Dean of Quality Assurance and Development Hayel University **KSA**.

#### Dr. Nawfal Jebbor

Department of Physics, Moulay Ismail University, Meknes, Morocco.

#### Dr. Eng.Ahmed Kadhim Hussein

Assistant Professor, Department of Mechanical Engineering, College of Engineering, University of Babylon, **Republic of Iraq.** 

#### Prof. Dr. Abd El Fady Beshara Morcos

Ass. Prof. of Relativistic Astrophysics and Cosmology, National Research In stitute of Astronomy and Geophysics, Egypt.

#### Zohre Bahrami

Shahid Beheshti University of Medical Sciences, Tehran, Iran. Researcher and Methodology Adviser.

#### Dr. Ayhan Kapusuzoglu

Department of Banking and Finance, Yildirim Beyazit University, Turkey.

#### **Dr. Charalambos Tsekeris**

Department of Psychology, Panteion University of Social and Political Sciences, Athens, Greece.

#### Dr. Mahdi Zowghi

Industrial and System Engineering, Management and Soft Computing, London Business and engineering School, **United Kingdom.** 

#### **Dr. Tomislav Jurendic**

Bioquanta Ltd. for Research and Development, Koprivnica, Croatia

#### Dr. Hanna Bolibok-Bragoszewska

Warsaw University of Life Sciences, Poland.

#### Prof. Md. Amin Uddin Mridha

Ph.D. DIC (London), Plant Production Department, King Saud University, P.O.Box 2460, Riyadh 11451, **Kingdom of Saudi Arabia** 

#### Dr. Alaa Abdelwahed Abdelbary

Prof. of Computational and Applied Mathematics, Arab Academy for Science and Technology & Maritime Transport, Egypt.

#### Dr. N R Birasal

Associate Professor, Zoology Department, KLE Society's G H College, HAVERI - 581 110, Karnataka state, India.

#### Dr. Nawab Ali Khan

Professor of Human Resource Management, College of Business Administration, Salman Bin Abdulaziz University, Post Box:165, Al Kharj - 11942 Kingdom of Saudi Arabia

#### **Editors**

#### Dr. Mukesh Kumar Meena

Assistant Professor (Crop Physiology), Department of Crop Physiology, University of Agricultural Sciences, Raichur-584104, Karnataka , India

#### Prof. Dr. Tarek Ahmed Shokeir

Professor and Consultant, Department of Obstetrics & Gynaecology, Fertility Care Unit, Mansoura University Teaching Hospitals, Mansoura Faculty of Medicine, **Egypt** 

#### Leila Falahati

Department of Resource Management and Consumer Studies, Faculty of Human Ecology, University Putra Malaysia.

#### Dr. Muhammad Ismail Mohmand

Tutor/Administrator in the Excellence Training Den College in Newcastle, United Kingdom

#### Noorbakhsh Hooti

Associate Professor in Dramatic Literature, Razi University, Faculty of Arts, English Department, Kermanshah, Iran

#### Dr. Ali Elnaeim Musa

University of Bahri, Sudan College of Applied and Industrial Sciences, Sudan

#### Prof. Dr. Magda M.A. Sabbour

Professor, Department of Pests and Plant Protection- National Research Centre, Cairo, Egypt.

#### Dr. Vahid Majazi Dalfard

Department of logistics and operation management, University of Vienna, Vienna, Austria

#### Dr. Basharia Abd Rub Alrasoul Abd Allah Yousef

Deputy Dean at Faculty of Engineering, University of Bahri, Khartoum, Sudan

#### Nasser Mousavi

Islamic Azad University, Bilesavar Branch, Iran.

#### Dr. Jinu John

Associate Professor (Biotechnology), Jinu Bhavan, Chepra (P. O), Kottarakara, Kollam (Dist.), Kerala - 691520; India.

#### Seyyed Mousa Hosseini

Head of Young Researchers Club, Islamic Azad University (Sama Organization), Mazandaranand, Gilan province, Iran.

#### Dr. Sunil Kumar

Assistant Professor, Department of Mathematics, National Institute of Technology, Jamshedpur, 831014, Jharkhand, India

#### Dr. Seyed Hossein Hosseini Nazhad

Department of Computer Engineering, Islamic Azad University, Iran

#### Zairi Ismael Rizman

Senior Lecturer, Faculty of Electrical Engineering, Universiti Teknologi MARA (UiTM) (Terengganu) Malaysia

#### Muhammad Attique Khan Shahid,

Associate Professor of Physics, Department of Physics, GC University, Faisalabad. **Pakistan.** PNRA certified Health Physicist, RPO, RSO Atomic and Nuclear Physics Lab

#### Mohsen Shafiei Nikabadi(PhD.)

Assistant Professor, Faculty of Economics and Management, Industrial Management Department, Semnan University, Semnan, Iran.

#### Dr.Vuda Sreenivasarao

Department of Computer and Information Technology, Defence University College, Deberzeit, Ethiopia

#### Dr. Mohdammed Israil

Post Doctoral Fellow, University Sains Malaysia, Pulau Penang, Malaysia.

#### Dr. S. Ravichandran

Assistant Professor, Department of Physics, Sathyabama University, India

#### Dr. Sukumar Senthil Kumar

School of Mathematical Sciences, Universiti Sains Malaysia, Malaysia.

#### Seifedine Kadry

American University of the Middle East, Kuwait.

#### Dr. Datta Asaram Dhale

Assistant Professor, Post Graduate Department of Botany, Ghogrey Science College, Dhule - Maharashtra State, India.

#### Dr. Ho Soon Min

Senior Lecturer, Faculty of Applied Sciences, INTI International University, Persiaran Perdana BBN, Putra Nilai, Negeri Sembilan, Malaysia.

#### Dr. Ezzat Molouk Kenawy

Economic Department, Faculty of Commerce, Kafr El-Sheikh University, Egypt.

#### Dr. Farooq Ahmad Gujar

Centre for Advanced Studies in Pure and Applied Mathematics, Bahauddin Zakariya University, Multan, 60800, **Pakistan.** & Head of Institution / Principal / Associate Professor of Mathematics.

#### Alireza Karbalaei

Assistant Professor, Department of English, Qeshm International Branch & Shariaty College, Iran.

#### Dr. Seshadri Sekhar. Tirumala

Principal, Chirala Engineering College, India.

#### Sayed Roholla Mousavi,

Department of Agriculture, Payame Noor University, Tehran, Iran.

#### Dr. Tarek Y. El-Hariri

Associated Professor, Egyptian Petroleum Research Institute, Exploration Department, Egypt.

#### Dr Mamode Khan Naushad

Department of Economics and Statistics, Faculty of social studies and humanities, University of Mauritius, Mauritius.

#### Dhahri Amel

Research professor, Research Unit: Materials, Energy and Renewable Energies (MEER)-Science Faculty of Gafsa, **Tunisia.** 

#### Dr. Muhammad Waqas Anwar

COMSATS Institute of Information Technology, University Road, 22060, Abbottabad, Pakistan.

#### Prof. Dr. Abdul-Kareem J.Al-Bermany

Advance Polymer Laboratory, Physics Department/College of Science/Babylon University, Iraq.

#### Dr. Bensafi Abd-El-Hamid

Assistant Professor, Dept. of Chemistry, Faculty of Sciences, Abou Bekr Belkaid University of Tlemcen, Algeria.

#### Dr. Vikas Anand Saharan

Assistant Professor & Head, Department of Pharmaceutics, Institute of Pharmaceutical Sciences & Drug Research, Seth GL Bihani SD College of Technical Education, **India**.

#### Dr. Syed Zulfiqar Ali Shah

Chairman Higher Studies and Research, Faculty of Management Sciences, International Islamic University Islamabad, **Pakistan.** 

#### Saima Anis Mustafa

Assistant Professor in COMSATS Institute of Information Technology, University Road, Abbottabad, Pakistan

#### Dr. Nagasamy Venkatesh

Assistant Professor, Dept. of Pharmaceutics, JSS College of Pharmacy, Tamil Nadu, India.

#### Mirza Hasanuzzaman

Department of Agronomy, Faculty of Agriculture, Sher-e-Bangla Agricultural University, Dhaka, Bangladesh.

#### Dr.K.V.L.N.ACHARYULU

Faculty of Science, Department of Mathematics, Bapatla Engineering college, Bapatla, India.

#### Maryam Ahmadian

Post Doctoral Fellow, Department of Social and Development Sciences, Faculty of Human Ecology, Universiti Putra, UPM Serdang, Selangor, Malaysia.

#### Dr. Mohammad Abul Hossain

Associate Professor, Department of Chemistry, University of Dhaka, Bangladesh.

#### Abdel Baset Hasoneh,

PhD, Associate professor of Marketing, Head of marketing Department Al Isra University - Amman, Jordan

#### Dr. Muhammad Akram

Faculty of Agriculture, Department of Eastern Medicine and Surgery, University of Poonch, Rawalakot, Azad Jamu and Kashmir, **Pakistan**.

#### **Dr. Anshoo Agarwal**

RAK Medical College and Health Sciences University, P.O.Box:13268, RAK, UAE, United Arab Emirates

#### Dr. Aamir Shhazad

Assistant Professor, Department of Physics, GC University, Faisalabad

#### Dr.(Mrs.) Sunanda Sharma

B.V.Sc & A.H., M.V.Sc., Ph.D. Department of Veterinary Gynecology and Obstetrics, College of Veterinary & Animal Science, Rajasthan University of Veterinary & Animal Sciences, Bikaner, **India**.

#### Muhamad Fazil bin Ahmad

Asst. Prof. Universiti Sultan Zainal Abidin, Terengganu, Malaysia.

#### **Shaukat Amer**

CPA, Assistant Professor, Department of Management Sciences, COMSATS Institute of Information Technology, Attock, **Pakistan.** 

#### Mohammad Hassan Boostani

Education Organization of Fars Province & Young Researchers and Elites Club, Zarghan Branch, Islamic Azad University, Zarghan, Iran

#### Naveed Ahmed

Assistant Professor, Department of business administration, Indus International Institute, 2-Km, Jampur Road, Dera Ghazi Khan, **Pakistan** 

#### **Rab Nawaz Lodhi**

PhD (ABD), Management Sciences (Bahria University Islamabad), Lecturer: Department of Management Sciences, COMSATS Institute of Information Technology, Sahiwal, **Pakistan.** International Licensed Trainer - NVivo Qualitative Research: QSR International Limited **Australia** 

#### Dr. Majid Sharifi Rad

Department of Range and Watershed Management, Faculty of Natural Resources, University of Zabol

#### Dr. Muhammad Naeem

LECTURER, Department of Information Technology, Hazara University, Mansehra.

#### Dr. Sohrab Mirsaeidi

Centre of Electrical Energy Systems (CEES), Faculty of Electrical Engineering (FKE), Universiti Teknologi Malaysia (UTM), 81310 Skudai, Johor, Malaysia

#### Dr. Muhammad Majid Gulzar

Faculty of Engineering, University of Central Punjab, Lahore, Pakistan

#### **Farhan Altaee**

Ministry of Science and Technology, Iraq-Baghdad

#### Dr. Hafiz Abdul Wahab

Assistant Professor of Mathematics, Department of Mathematics, Hazara University Mansehra Pakistan.

#### **Dr. Rohit Bansal**

Assistant Professor, Department of Management Studies, Vaish College of Engineering, Rohtak (Haryana), India

#### Dr. Muhammad Akram

Faculty of Agriculture, Department of Eastern Medicine and Surgery, University of Poonch, Rawalakot, Azad Jamu and Kashmir, **Pakistan**.

#### Dr. Meena M.K.

M.Sc.(Agri.), Ph.D., Assistant Professor, Department of Crop Physiology, University of Agricultural Sciences, Raichur-584104,Karnataka, **India**.

## **Table of Contents, November 2022**

Hala Kandil, A. Khater, A. El-Hassanin and Amal El-Maghraby

Using Nutrient Elements, Humic Acids and Amino Acids to Maximize Tomato Plants Production under Soil Salinity Conditions: 1. Effect on Growth and Macronutrient Status

J. Basic Appl. Sci. Res. 2022 12(2): 1-14. [Abstract] [Full-Text PDF]

\_\_\_\_\_



*J. Basic. Appl. Sci. Res.*, 12(2)1-14, 2022 © 2022, TextRoad Publication ISSN 2090-4304 Journal of Basic and Applied Scientific Research www.textroad.com

## Using Nutrient Elements, Humic Acids and Amino Acids to Maximize Tomato Plants Production under Soil Salinity Conditions: 1. Effect on Growth and Macronutrient Status.

Hala Kandil<sup>1</sup>, A. Khater<sup>1</sup>, A. El-Hassanin<sup>2</sup> and Amal El-Maghraby<sup>1</sup>

<sup>1</sup>Plant Nutrition Department, National Research Centre, 33 El Buhouth St., Dokki, Cairo 12622, Egypt. <sup>2</sup>Dept.of Natural Resources, Faculty of African Postgraduate Studies, Cairo University, Egypt.

> Received: July 13, 2022 Accepted: October 8, 2022

#### ABSTRACT

A pot experiments were conducted at greenhouse of National Research Center. using sample collected from El Husseineya - Port Said, Egypt. Experiment was carried out to studies the best soil addition treatments maximized growth of tomato plants under soil salinity conditions. The treatments are: (Control, Humic acid 2%, Algae extract 2%, Potassium 500 ppm, Silica 8 mM, Proline 100 ppm).

Addition of humic acid increase percent of plant height and root length by 29.34%, 38.06% then potassium by 26.73%, 27.97%, silica by 23.65%, 14.7%, algae 20.36%, 9.96% and finally proline 14.58%, 8.04%. Also, humic acid treatment gave best growth (fresh and dry weight) of tomato plants while the lowest one was proline treatment.

All soil additions give significant positive effect on N content compared with control, the increase ranged between 2.47% up to 4.51% in shoot and from 1.38% up to 3.22% in root. The maximum values of P content obtained using humic acid 0.44%, 0.33% for shoot and root coppered with control. The highest K concentration values in shoot and root of tomato plants were obtained by using K treatment (6.35%, 5.94%) while the lowest ones were attained by applying proline treatment (3.04%, 2.47%) for shoot and root respectively.

Generally, the best tretments were humic acid, potassium and algae which gave a significant increase in all the growth parameters and helped tomato plants grown in saline soil condition.

KEY WORDS: Algae, Humic acid, organic matter, Potassium, Proline, Salinity, Silica, Tomato.

#### Introduction

Tomato (Lycopersicon esculentum Mill.), belonging to the family Solanaceae, is one of the most important, popular, nutritious and palatable vegetables Egypt is one of the major tomato producing and exporting country. Most land is cropped at least twice a year, but agricultural productivity is limited by salinity, which afflicts an estimated 35% of cultivated land, and drainage problems. Tomato is one of the most important, popular, nutritious and palatable vegetables grown in Egypt. It plays a vital role in providing a remarkable quantity of vitamin A and vitamin-C in human diet. It can be eaten both in raw as well as ripe and after cooking. The production of tomatoes in Egypt was approximately 7 million metric Tons.

Corresponding Author: Hala Kandil, Plant Nutrition Department, National Research Centre, 33 El Buhouth St., Dokki, Cairo 12622, Egypt Citation: Hala Kandil, A. Khater, A. El-Hassanin and Amal El-Maghraby; 2022, Using Nutrient Elements, Humic Acids and Amino Acids to Maximize Tomato Plants Production under Soil Salinity Conditions: 1. Effect on Growth and Macronutrient Status.; Journal of Basic and Applied Scientific Research, 12(2)1-14.

Salinity in the soil causes limitation in the plant growth, productivity particularly in arid and semi-arid areas (Shannon, 1998). Salinity is the level of salt dissolved in water. Salinity can refer to the salt contained in the soil. Saline conditions are often found in lowland areas. High concentrations of salt can cause high amounts of  $Na^+$  to accumulate easily in plant cells, which is toxic at high concentrations (Mathius, 2014).

**Gupta**, (1987) found that the organic and inorganic amendments are effective in the amelioration of saline soils. Organic matter decomposition and plant root action also help dissolve the calcium compounds found in most soils, thus promoting reclamation of saline soil.

The addition of organic matter or its extracts to the soil has a very important effect on the various physical, chemical, and biological properties of the soil, as it is an important source of many nutrients and an essential source of energy for soil organisms (Tan, 2014; Al-Khafaji, 2015 and Abd El-Monem et al., 2008). Some organic extracts (chelates) such as humic acid have a significant role in improving plant growth and increasing the efficiency of roots in absorbing water and dissolved nutrients, increasing the plant's ability to tolerate salt stress, and improving various physical, chemical, and biological properties of the soil (Jasim et al., 2015 and Ragab et al., 2008).

Potassium is one of the critical macro-elements for plant growth because it is involved in some physiological processes such as (1) biophysical processes in its role in controlling osmotic, cell turgor, pH stability, and (2) biochemical processes in its role in the activity of enzymes in synthesizing carbohydrates and proteins as well as increasing photosynthetic translocation (Amisnaipa et al., 2009). Adding K to the soil can neutralize the land from Na stress, and this works for some plants, including tomatoes (Pujiasmanto et al., 2010).

The beneficial effects of silicon (Si) are mainly linked with high deposition in plant tissues, enhance its strength and stiffness, this increase led to mechanical strength reduces housing, pest attacks and increases the light receiving position of the plant, increase photosynthesis and promote growth (Epstein, 1999), also supposed to reduce transpiration from the cuticle thus increasing resistance to salinity (Matoh et al., 1986).

Proline is known to accumulate under conditions of environmental stresses to play a role in the process of osmotic adjustment in many crops (Heuer, 1994). Their main role is probably to protect plant cells against the ravages of salt by preserving the osmotic balance (Heuer, 2003, Ashraf and Foolad, 2007).

Algae have been used for the production of plant biostimulants, i.e. of bioproducts which contain substances "whose function, when applied to plants or the rhizosphere, is to stimulate natural processes, to enhance/benefit nutrient uptake, nutrient efficiency, tolerance to abiotic stress, and crop quality". (European Biostimulant Industry Council, EBIC 2014). Algae contain important quantities of plant growth hormones, auxins, abscisic acid, cytokinins, gibberellins (Tarakhovskaya et al., 2007).

The aim of this investigation was evaluated of effect of some treatments for reducing salinity damage on Growth, Yield and Quality of tomato.

#### MATERIALS AND METHODS

Greenhouse experiment was conducted using saline soil to study the impact of some soil treatments (element nutrition and amino acid) to minimize adverse effect of salinity on plant growth.

Surface soil sample was collected from El Husseineya - Port Said, Egypt. Before being used, the soil samples were air dried, pulverized, sieved through a sieve 2mm, and thoroughly combined. The physicchemical characteristics of the studied soil were determined according to international methods and summarized in Table (1).

Physical properties										
Coarse sand	Fine sand	l Silt	Cla	ay	Texture	Field ca	pacity	Wilting point	Ava w	ailable ater
%										
6.73	23.90	30.27	39.10	10 Clay loam 32.		32.43		13.44	18.99	)
Chemical properties										
pН	EC	O.M	.M CaCO <sub>3</sub> Macronutrients (mgkg <sup>-1</sup> )							
(1:2.5)	(dSm <sup>-1</sup> )	(%	6)		Ν		Р		К	
8.12	9.88	0.58	11.75	34.60			4	1.22	187	.00
Cations (meql <sup>-1</sup> )				A	Anions (meql <sup>-1</sup> ) Micronutrients (mgkg <sup>-1</sup> )					
Ca <sup>++</sup>	Mg <sup>++</sup>	Na <sup>+</sup>	<b>K</b> <sup>+</sup>	HCO3 <sup>-</sup>	Cl	SO <sup>=4</sup>	Fe	Mn	Zn	Cu
12.55	22.10	63.34	0.81	9.77	57.20	31.83	6.12	2.88	0.59	1.18

Table (1): Some of the soil's physical and chemical features.

Plastic pots with dimensions (20\* 24 cm) were used. They were filled with 5 kg of soil that was mixed and homogeneous. The recommended fertilizer treatments for each pot were placed before planting, which are (ammonium sulfate 300 kgfed<sup>-1</sup>) - potassium sulfate 200 kgfed<sup>-1</sup> superphosphate 100 kgfed<sup>-1</sup>500 ml l.

Tomato seedlings of hybrid Alicia variety, which were get from the Agricultural Research Center in El Husseineya, Port Said, planted on April 16<sup>th</sup>, by placing two seedlings in each pot, irrigated with 70% of field capacity and maintain the moisture level by irrigation every 4 days.

After two weeks from transplanting, on May 2, the treatments have been applied with 4 replicate for each treatment.

The treatments are:500 ml l for every pot from soluatian of water as control, Humic acid 2g l<sup>-1</sup>, Algae extract 2%, Potassium 500 ppm as potassium nitrate, Silica 8 mM (as calcium silicate andProline 100 ppm

Citation: Hala Kandil, A. Khater, A. El-Hassanin and Amal El-Maghraby; 2022, Using Nutrient Elements, Humic Acids and Amino Acids to Maximize Tomato Plants Production under Soil Salinity Conditions: 1. Effect on Growth and Macronutrient Status.; Journal of Basic and Applied Scientific Research, 12(2)1-14.





General photos for pot experiment.

All of the pots are placed in a randomized block pattern. On May 17<sup>th</sup>, at the vegetative development stage, chlorophyll was detected in the leaves of tomato plants.

At the beginning of the flowering stage( June 24) plants were collected, after 68 days of transplanting, and deferent parameters were recorded (plant length - branches number per plant - leaves number per plant - fresh and dry weight) and determents (N, P, K, Na, Fe, Mn, Zn and Cu).

Analytical statistics: The findings were analyzed using Snedecor and Cochran's (1982) analysis of variance, and the treatments were compared using the L.S.D test at the 0.05 levels of probability.

#### **RESULTS AND DISCUSSION**

The experiment aims to apply some soil treatments that previous studies have referred to, with regard to their ability to reduce the harmful effects of salt on tomato plants, to choosing best treatment. Therefore, humic acid (H.A), algae, potassium as KNO<sub>3</sub>, silica and proline were employed to investigate their impact on the development of tomato plants in a saline soil from El Husseineya area. To study its effect on the vegetative characteristics of tomato plants and the nutritional content under the influence of saline soil.

In many semi-arid locations, such as the Mediterranean, where salt has been a major limit in agricultural production, the tomato plant is regarded a significant greenhouse crop.

#### Vegetative growth:

Data presented in Table (2) and illustrated in Fig (1, 2) show the effect of soil treatments on plant height and root length of tomato plants. All treatment used are given significant increases compared with control. These increases can be attributed to improving effect of soil additions to help tomato plants for growth under salinity condition. The obtained data agree with those found by **Khaled and Fawy (2011)**, **Hala Kandil et al., (2020)**, **Chakraborty, et al., (2016)**, **Ali et al., (2012) and Tala et al., (2013)** who found that addition humic acid, algae, potassium, silica and proline increased all growth parameters for corn, parsley, peanut and wheat respectively.

Data revealed that addition of humic acid increased plant height and root length by 29.34%, 38.06% followed by potassium 26.73%, 27.97% then silica by 23.65%, 14.7% then algae 20.36%, 9.96% and finally proline 14.58%, 8.04%. On the other hand leaf number gives non- significant increase with algae, silica and proline treatment compared with control.

The highest plant height, root length and leaf number recorded under humic acid treatment (43.2 cm, 11.5 cm and 13.66) respectively. This confirm with results by **Türkmen et al.**, (2004) and Paksoy et al., (2010) who reported that the humic acid treatment improved the growth characteristics of plants growing in a saline environment. Khaled and Fawy (2011) also discovered that the most cost-effective treatment amounts should be selected and should not exceed 2 g humus kg<sup>-1</sup> in soil and 0.1 percent in foliar treatment.

Husseineya soil.							
Treatments	plant height	Root length	Leaf number per plant				
	(cm)	(cm)					
Control	33.40 <sup>f</sup>	8.33 <sup>f</sup>	10.33 <sup>b</sup>				
H.A	43.20ª	11.50ª	13.66ª				
Algae	40.20 <sup>d</sup>	9.16 <sup>d</sup>	11.33 <sup>b</sup>				
K	42.33 <sup>b</sup>	10.66 <sup>b</sup>	13.66 <sup>ab</sup>				
Si	41.30°	9.56°	11.33 <sup>b</sup>				
Proline	38.27°	9.00 <sup>e</sup>	11.66 <sup>b</sup>				
LSD	0.183	0.074	2.179				

 Table (2): Effect of some soil application on some growth parameters for tomato plant in El

 Husseineva soil.

Citation: Hala Kandil, A. Khater, A. El-Hassanin and Amal El-Maghraby; 2022, Using Nutrient Elements, Humic Acids and Amino Acids to Maximize Tomato Plants Production under Soil Salinity Conditions: 1. Effect on Growth and Macronutrient Status.; Journal of Basic and Applied Scientific Research, 12(2)1-14.



Fig. (1): Effect of some soil application on plant height and root length for tomato plant in El Husseineya soil.



Fig. (2): Effect of some soil application on leaf number for tomato plant in El Husseineya soil.

Data of Fresh and dry weights are presented in table (3) and illustrated in figs (3 and 4). The obtained data indicated that fresh and dry weights of tomato plants grown under saline soil give positive response for all soil additions compared with control.

Data revealed an increasing in dry weight percentage of shoot and root tomato plants for humic acid (68% and 71.4%) these results confirm with **Ouni et al.**, (2014) who reported that the direct actions for humic acid effects on germination, plant growth (root and shoot), and hormone-like activity on plants are attributable to their effects on germination.

In general, H.A treatment gave best growth (fresh and dry weight) of tomato plants and the lowest growth was attained under proline treatment as compared to other forms of therapy. These findings are consistent with previous findings by **Cimrin et al. (2010)** they found that Pepper plant growth was improved by humic acid treatment in terms of fresh and dry weight, shoot and root lengths, shoot breadth, cotyledon length and breadth, and hypocotyls length.

Treatment	Fresh v	veight (g)	Dry wei	ight (g)
	Shoot	Root	Shoot	Root
Control	7.63 <sup>f</sup>	0.93°	2.50 <sup>d</sup>	0.35 <sup>d</sup>
H.A	12.76ª	1.80ª	4.20ª	0.60ª
Algae	10.13 <sup>d</sup>	1.53 <sup>b</sup>	3.30 <sup>bc</sup>	0.53 <sup>b</sup>
K	11.40 <sup>b</sup>	1.31°	3.75 <sup>ab</sup>	0.50 <sup>b</sup>
Si	10.63°	1.23 <sup>d</sup>	3.50 <sup>a-c</sup>	0.43°
Proline	9.36°	0.83 <sup>f</sup>	3.00 <sup>cd</sup>	0.43°
LSD	0.127	0.0178	0.723	0.032

Table (3): Effect of soil treatments on Fresh and dry weight of tomato plant



Fig. (3): Effect of soil treatments on fresh weight (shoot and root) for tomato plant



Fig. (4): Effect of soil treatments on dry weight (shoot and root) for tomato plant

#### Macronutrients:

Nitrogen (N), Phosphorus (P) and Potassium (K<sup>+</sup>) are key macronutrients that have a role in a variety of plant processes, particularly storage and energy transport. Plant development requires soil macronutrients (N, P, and K).

Nutrients play a variety of roles in the structure, metabolism, and osmoregulation of plant cells in general (Taiz and Zeiger, 2009). Nutritional problems, which are caused by salinity's influence on the availability, absorption, and transport inside the plant, are one of the most significant salt stress impacts on plants (Munns and Tester, 2008). The negative effects of salinity on plant development and production are ascribed by nutrient shortage, ion toxicity, and osmotic stress (Nublat et al., 2001).

Citation: Hala Kandil, A. Khater, A. El-Hassanin and Amal El-Maghraby; 2022, Using Nutrient Elements, Humic Acids and Amino Acids to Maximize Tomato Plants Production under Soil Salinity Conditions: 1. Effect on Growth and Macronutrient Status.; Journal of Basic and Applied Scientific Research, 12(2)1-14.

Because more Na and Cl ions compete with nutrients like  $K^+$ ,  $Ca^{2+}$ , and  $NO_3^-$ , salinity can create nutrition deficits or imbalances. On the other side, it has the potential to influence nutrient absorption and hinder nutrient translocation. Improved fertilizer management in arid and semiarid settings by a better knowledge of the function of mineral nutrients in plant tolerance to salt (**Hu et al., 2017**).

This part of the experiment was concerned with the effect of soil treatments on tomato plants' macronutrient (N-P-K).

#### Nitrogen (N):

Nitrogen makes for around 80% of total mineral nutrients. Its availability is the most extensively limiting factor for plant growth and productivity under salt stress. It has been discovered that the interaction between salinity and nitrogen occurs more frequently in soils that are low in nitrogen. Many researches have shown that salinity reduces nitrate content in plants while having little effect on total nitrogen content, and this might be the situation in this study (Hu and Schmidhalter 2005).

Data in table (4) and illustrated in Fig. (5) revealed that nitrogen content in shoot and root of tomato plants were significantly increased affected by all treatments as compared with control the increase ranged between 2.47% up to 4.51% in shoot and from 1.38% up to 3.22% in root. Nitrogen contents were higher in shoot than in root for all treatments.

These results harmony with **Khaled and Fawy (2011)** who found that the corn absorption of nitrogen was enhanced when humus was applied to the soil, **Sabagh et al. (2017)** Explained that the effect of proline under saline conditions causes an increase in the nodule number and Nitrogen Fixation .Also, when proline was added, it was discovered that the loss of nitrogenase activity caused by salt could be overcome. **El-Lethy et al., (2013)** found that the potassium application, especially at 150 mg kg<sup>-1</sup> soil had significantly caused increases in macro elements.

Humic acid gave the highest percentage of increase N compared to control (82.6%, 133%) for shoot and root and the lowest percentage increase values obtained at proline treatments (12.6%, 36.9%) for shoot and root respectively.

**Turkmen et al. (2004)** found that tomato grown on a saline medium supplemented with humic acid had a considerable improvement in mineral nutrient absorption. This is most likely related to the increased permeability of root cell membranes. The humic chemical may also function by promoting root development and proliferation. These effects are especially crucial for plant adaptation to harsh soil conditions like salinity.

nussenicya son.								
	treatments	N (%)		P (%	6)	K (%)		
		Shoot	Root	Shoot	Root	Shoot	Root	
	Control	2.47°	1.38 <sup>f</sup>	0.2 <sup>f</sup>	0.18 <sup>d</sup>	3.18 <sup>e</sup>	1.91 <sup>f</sup>	
	H.A	4.51 <sup>a</sup>	3.22 <sup>a</sup>	0.44 <sup>a</sup>	0.33ª	5.17 <sup>b</sup>	4.35 <sup>b</sup>	
	Algae	3.45 <sup>d</sup>	2.15 <sup>d</sup>	0.36°	0.27 <sup>b</sup>	4.02 <sup>d</sup>	3.67°	
	K	3.93ª	2.82 <sup>b</sup>	0.38 <sup>b</sup>	0.29 <sup>b</sup>	6.35 <sup>a</sup>	5.94ª	
	Si	3.19 <sup>e</sup>	2.77°	0.31 <sup>d</sup>	0.21°	4.49°	3.11 <sup>d</sup>	
	Proline	2.78 <sup>f</sup>	1.89 <sup>e</sup>	0.25 <sup>e</sup>	0.18 <sup>d</sup>	3.04 <sup>f</sup>	2.47 <sup>e</sup>	
	LSD	0.178	0.252	0.017	0.025	0.177	0.184	

 Table (4): Effect of some soil addition on macronutrients content for tomato plant in El

 Husseineya soil.

Kandil et al., 2022



Fig. (5): Effect of some soil addition on nitrogen content for tomato plant

#### **Phosphorus:**

Depending on the plant type, growth stage, salt, degree of salinity, and P content throughout development, the interaction of phosphorus (P) with salt in plant nutrition is quite complicated and often confusing (Ashraf et al., 2009).

Phosphate absorption and buildup in soil-grown crops is hampered by salinity, which limits phosphate availability (**Prasad et al., 2014**).

Because of ionic strength effects, sorption processes in soil solution, and low solubility of Ca-P minerals in salty soils, mineral nutrients such as P availability is reduced.

Data in table (4) and illustrated in Fig. (6) revealed that significant differences in P content in shoot and root of tomato plant. Phosphorus content value in shoot under all treatments higher than P content value in root under the same treatment. The maximum increase values of P content obtained using humic acid (0.44%, 0.33%) for shoot and root compared with control, similar results obtained by **Turkmen et al. (2004)**,

In case of potassium treatment et was (0.38%, 0.29%) these results confirmed with El-Lethy et al., (2013) who found that potassium application, especially at 150 mg/kg soil had significantly caused increases in macro elements (N and P), in case of algae et was (0.36%, 0.27%) these results agree with Atzori et al., (2020) who found that algae application showed high P uptake efficiency, in case of silica et wase (0.31%, 0.21%) these results are in line with Murillo-Amador *et al.*, (2007) who improved cowpea and kidney bean growth, physiological parameters, balanced nutrition, and nutrient intake. Final proline case an increase with(0.25%, 0.18%) these results agree with Exogenous proline, according to Hossain and Fujita (2010), protects against salt-induced oxidative damage by lowering H2O2 and lipid peroxidation levels while also improving antioxidant defence and methylglyoxal detoxification systems.



Fig. (6): Effect of some soil addition on phosphorus content for tomato plant in El Husseineya soil.

Citation: Hala Kandil, A. Khater, A. El-Hassanin and Amal El-Maghraby; 2022, Using Nutrient Elements, Humic Acids and Amino Acids to Maximize Tomato Plants Production under Soil Salinity Conditions: 1. Effect on Growth and Macronutrient Status.; Journal of Basic and Applied Scientific Research, 12(2)1-14.

#### Potassium:

Potassium is an essential macronutrient for plant growth, development, production, and quality, as well as for plant survival under abiotic stress, as stress disrupts plant physiological systems, root and shoot elongation, enzyme activity, water and assimilate transfer, protein synthesis, photosynthetic transport, and chlorophyll content are all negatively affected by stress (Véry and Sentenac, 2003; Pettigrew, 2008 and Gerardeaux et al., 2010). Because potassium competes with sodium for binding and maintaining plant water status, it helps for salinity tolerance (Capula-Rodrguez et al., 2016).

The influence of different treatments on potassium content of shoot and root of tomato plants grown on El-Husseniya soil are presented in Table (4) and illustrated in Fig. (7). Data show that the application of different treatments when compared to control, generated a considerable rise in K concentration in the shoot and root of tomato plants and potassium concentration in shoot is bigger than root under the same treatments.

The highest K concentration values in shoot and root of tomato plants were obtained by using K treatment (6.35%, 5.94%) for shoot and root respectively. While the lowest ones were attained by applying proline treatment (3.04%, 2.47%) for shoot and root respectively.

Plants suffer from K shortage when they are exposed to high salt. As a result, boosting the Knutritional status of plants under salinity stress mitigates the negative impacts of Na<sup>+</sup> through a variety of mechanisms, including  $K^+ = Na^+$  discrimination (**Rodriguez-Navarro, 2000; Rubio et al., 2009)**. Furthermore, (**Shabala and Pottosin, 2014**) discovered that high  $K^+$  availability improves plant development under salt stress, but low  $K^+$  salt stress is harmful.



Fig. (7): Effect of some soil addition on potassium content for tomato plant in El Husseineya soil.

#### REFRENCES

- Abd El-Monem, E.A.A.; Saleh, M.M.S.; Mostafa, E.A.M. (2008). Minimizing the quantity of mineral nitrogen fertilizers on grapevine by using humic acid, organic and biofertilizers. Res. J. Agric. Biol. Sci., 4, 46–50.
- Ali, A.; Iqbal, N.; Ali, F.; Afzal, B.; Alternanthera, B. and Regel, G. (2012). Nicholson, a potential halophytic ornamental plant: Growth and physiological adaptations. Flora, 207: 318–321.

- Al-Khafaji, M. (2015). Effect of Concentration and spraying date of Humic acid on Growth and yield of Zea mays L. Kufa J. Agric. Sci., 7, 156–170.
- Amisnaipa, A.D. Susila, R. Situmorang, D.W. Purnomo. (2009). Penentuan Kebutuhan Pupuk Kalium Untuk Budidaya Tomat Menggunakan Irigasi Tetes Dan Mulsa Polyethylen. J. Agron. Indonesia.
- Ashraf, M. and Foolad, M.R. (2007). Roles of glycine betaine and proline in improving plant abiotic stress resistance. Environ. Exp. Bot. 59: 206-216.
- Ashraf, M.; Ozturk, M. and Athar, H. (2009). Salinity and Water Stress: Improving Crop Efficiency. Tasks for Vegetation Science Series,44 th Edition Springer Nature. by M. Ashraf (Editor), M. Ozturk (Editor), Habib-ur-Rehman Athar (Editor).
- Atzori, G.; Nissim, W.G.; Rodolfi, L.; Niccolai1, A.; Biondi1, N.; Mancuso, S. and Tredici, M.R. (2020). Algae and Bioguano as promising source of organic fertilizers. Journal of Applied Phycology, 32:3971–3981.
- Black, C.A.; D.D. Evans; L.E. Ensminger; G.L. White and F.E. Clarck (1982). 'Methods of Soil Analysis', Part 2. Agron. Inc. Madison Wise.
- Capula-Rodríguez, R.; Valdez-Aguilar, L.A.; Cartmill, D.L.; Cartmill, A.D. and Alia-Tejacal, I. (2016). Supplementary calcium and potassium improve the response of tomato (*Solanum lycopersicum L.*) to simultaneous alkalinity, salinity, and boron stress. Commun. Soil Sci. Plant Anal., 47 (4): 505–511
- Chakraborty, K.; Bhaduria, D.; Meena, H.N. and Kalariya, K. (2016). External potassium (Kþ) application improves salinity tolerance by promoting Nab-exclusion, Kb-accumulation and osmotic adjustment in contrasting peanut cultivars. Plant Physiology and Biochemistry, 103: 143-153.
- Cottenie, A.; Verlea, M.; Krekens, L.; Velghe, G. and Camerlynck, R. (1982). Chemical Analysis of Plant and Soils Lab. Anal. Agroch. Fac. Agric. State University Gent., Belgium. 1-63.
- Çimrin, K.M., Türkmen, Ö., Turan, M., Tuncer, B., (2010). Phosphorus and humic acid application alleviate salinity stres of pepper seedling. African Journal of Biotechnology, 9 (36): 5845-5851.
- **EBIC** (European biostimulants industry council), (2014). Promoting the biostimulant industry and the rule of plant biostimulants in making agriculture more sustainable. (http://www.biostimulants.eu/).
- El-Lethy, S.R.; Magdi, T.A. and Reda, F. (2013). Effect of Potassium Application on Wheat (Triticum aestivum L.)Cultivars Grown Under Salinity Stress. World Applied Sciences Journal, 26 (7): 840-850.

- **Epstein, E. (1999)**. Silicon: Annual Review of Plant Physiology and Plant Molecular Biology, 50, 641-664.
- Gerardeaux, E.; Jordan-Meille, L.; Constantin, J.; Pellerin, S. and Dingkuhn, M. (2010). Changes in plant morphology and dry matter partitioning caused by potassium deficiency in Gossypium hirsutum (L.). Environ Exp Bot 67:451–459.
- Gupta SK, Gupta IC. (1987). Management of Saline Soils and Waters. Oxford and IBH Publication, New Delhi.
- Hala Kandil; Mahmoud, S.A.; Siam, H. and Abd El-Aty, A.M. (2020). Response of Parsley (*Petroselinum Crispum*) to microalgae Chlorella Vulgaris and Anabaena Sphaerica as biofertilizers. Plant Archives, 20 (2): 9561-9567.
- Heuer, B. (1994). Osmoregulatory role of proline in water and salt stressed plants. In: Pessarakli, M. (Ed.), Handbook of Plant and Crop Stress. Marcel Dekker, New York, pp. 363–381.
- Heuer, B. (2003). Influence of Exogenous Application of Proline and Glycinebetaine on Growth of Salt-Stressed Tomato Plants. Plant Science, 165, 693-699.
  Cimrin, K.M., Turkmen, O., Turan, M. and Tuncer, B. (2010). Phosphourus and humic acid application alleviate salinity stress ofbpepper seedling. African Journal of Biotechnology, 9(36): 5845-5851.
- Hossain, M.A. and Fujita, M. (2010). Evidence for a role of exogenous glycinebetaine and proline in antioxidant defense and methylglyoxal detoxification systems in mung bean seedlings under salt stress. Physiol. Mol. Biol. Plants, 16(1): 19-29.
- Hu, W.; Coomer, T.D.; Loka, D.A.; Oosterhuis, D.M. and Zhou, Z. (2017). Potassium deficiency affects the carbon-nitrogen balance in cotton leaves. Plant Physiol. Biochem. 115: 408–417.
- Hu, Y. and Schmidhalter, U. (2005). Drought and salinity: a comparison of their effects on mineral nutrition of plants. Journal of Plant Nutrition and Soil Science, 168: 541-549.
- Jackson, M.L. (1973). Soil Chemical Analysis. Prentice-Hall, Inc., Englewood Cliffs, USA.
- Jasim, A.H.; Alryahii, I.; Abed, H.M.; Badry, A.N. (2015). Estimation Effect of some treatments on alleviating of environmental stress on growth and yield of squash (*cucurbittapepo L.*). Mesop. Environ. J. 1, 67–74.
- Khaled, H. and Fawy, H.A. (2011). Effect of different levels of humic acids on the nutrient content, plant growth, and soil properties under conditions of salinity. Soil & Water Res., 6(1): 21-29.
- Klute, A. (1986). Methods of soil analysis. Part 1. 2 nd Edition, Agronomy Series No. 9. Soil Sci. Soc. Am. Inc., Medison, WI, USA.

- Lindsay, W.L. and Norvell, W.A. (1978). Development of DTPA soil test for zinc, iron, manganese and copper. Soil Sci. Soc. Amer. J., 42: 421-428.
- Mathius, F.J.M. (2014). Sodium in plants: perception, signalling, and regulation of sodium fluxes. Journal of Experimental Botany 66 (3): 849-858.
- Matoh, T., Kairusmee, P. and Takahashi, E. (1986). Salt induced damage to rice plants. Australian J. Basic and Applied Sci., 5 (9): 769-781.
- Munns, R. and Tester, M. (2008). Mechanisms of Salinity Tolerance. Annual Review of Plant Biology59:651-681.
- Murillo-Amador, B.; Yamada, S.; Yamaguchi, T.; Rueda-Puente, E.; Avila-Serrano, N; Garcia-Hernandez, J.L.; Lopez-Aguilar, R.; Troyo-Dieguez, E. and Nieto-Garibay, A. (2007). Influence of calcium silicate on growth, physiological parameters nutrition in two legume species under salt stress. J. Agronomy & Crop Science, 193: 413- 421.
- Nublat, A.; Desplans, J.; Casse, F. and Berthomieu, P. (2001). sas1,an Arabidopsis mutant overaccumulating sodium in the shoot, shows deficiency in the control of the root radial transport of sodium. The Plant Cell, 13:125-137.
- Olsen, S.R.; Cole, C.V.; Watnabe, F.S. and Dean, L.A. (1954). Estimation of available phosphorus in soil by sodium bicarbonate. U. S. Dept. Agric. Circ., 399.
- Ouni, Y.; Ghnaya, T.; Montemurro, F.; Abdelly, Ch. and Lakhdar, A. (2014). The role of humic substances in mitigating the harmful effects of soil salinity and improve plant productivity. International Journal of Plant Production, 8(3): 353-374.
- Paksoy, M.; Türkmen O. and Dursun A. (2010). Effects of potassium and humic acid on emergence, growth and nutrient contents of okra (*Abelmoschus esculentus* L.) seedling under saline soil conditions. African Journal of Biotechnology, 9(33): 5343-5346.
- Pettigrew, W.T. (2008). Potassium influences on yield and quality production for maize, wheat, soybean and cotton. Physiol Plant 133:670–681.
- Prasad, S.M.; Parihar, P. and Singh, V.P. (2014). Effect of Salt Stress on Nutritional Value of Vegetables. Biochemistry & Pharmacology, 3(2):1-2.
- Pujiasmanto, B., Sumiyati, H. Widijanto, and Alfiatun, A. (2010). Evaluation of giving legin and K fertilizer on the growth of soybean (*Glycine max (L.) Merrill*) under NaCl stress. Sains Tanah; J. Ilmu Tanah dan Agroklimatologi 7 (1): 16-24 (in Indonesian).
- Ragab, A.A.M.; Hellal, F.A.; Abd El-Hady, M. (2008). Water salinity impacts on some soil properties and nutrients uptake by wheat plants in sandy and calcareous soil. Aust. J. Basic Appl. Sci., 2, 225–233.

- **Rodriguez-Navarro, A. (2000).** Potassium transport in fungi and plants. Biochim Biophys Acta Rev Biomembr., 1469: 1–30.
- Rubio, J.S.; Garcia-Sanchez, F.; Rubio, F. and Martinez, V. (2009). Yield, blossom-end rot incidence, and fruit quality in pepper plants under moderate salinity are affected by K+ and Ca2+ fertilization. Sci Hortic-Amsterdam, 119: 79–87.
- Sabagh, A.EL.; Sorour, S.; Ragab, A.; Saneoka, H. and Islam, M.S. (2017). The Effect of Exogenous Application of Proline and Glycine Betaineon the Nodule Activity of Soybean Under Saline Condition. Journal of Agriculture Biotechnology,2(1): 01-05.
- Shabala, S. and Pottosin, I. (2014). Regulation of potassium transport in plants under hostile conditions: implications for abiotic and biotic stress tolerance. Physiol. Plant, 151, 257–279.
- Shannon, M.C. (1998). Adaptation of plants to salinity. Adv. Agron. 60, 75-119.
- Snedecor, G. and Cochran, W. (1982). Statistical methods. 6<sup>th</sup> Ed. Iowa state Univ. Press. Amer. Iowa, USA.
- Taiz, L. and Zeiger, E. (2009). Fisiologia Vegetal. 4th ed. Porto Alegre: Editora ARTMED. 848 p.
- Tala A.; Nawaz, K.; Hussian, K. and Bhatti, K.H. (2013). Foliar application of proline for salt tolerance of two wheat (*Triticum aestivum L.*) cultivars. World Applied Sciences Journal, 22(4):547-554.
- Tan, K.H. (2014). Humic Matter in Soil and the Environment: Principles and Controversies; CRC Press Taylor & Francis Group: Boca Raton, FL, USA.
- Tarakhovskaya, E.R., Maslov, Y.I., Shishova, M.F. (2007). Phytohormones in algae. Russian Journal of Plant Physiology, 54, 163–170.
- Türkmen, O.; Dursun, A.; Turan, M. and Erdinc, C. (2004). Calcium and Humic Acid Affect Seed Germination, Growth and Nutrient Content of Tomato (*Lycopersicon esculentum* L.) Seedlings under Saline Soil Conditions. Soil Plant Sci. 54: 168-174.
- Véry, A.A and Sentenac, H. (2003). Molecular Mechanisms and Regulation of K<sup>+</sup> Transport in Higher Plants. Annual Review of Plant Biology 54(1): 575-603.

## **INSTRUCTION TO AUTHORS**

### **Manuscript Submission:**

Send your manuscript with attachment by mailing it to <u>info@textroad.com</u>, <u>textroadjournals@gmail.com</u> along with <u>covering letter</u>.

#### **Manuscript Preparation:**

- \* Title
- \* Author names and addresses
- \* Abstracts (Not more than 300 words)
- \* Key words
- \* Introduction
- \* Materials and Methods
- \* Results and Discussions
- \* References (Use numbering in the text instead of full references). Give full references at the end of the file
- \* Photographs should be of high quality (Minimum 300-600 dpi)
- \* Graphs should be in clearly visible form so that it may become easy to redraw
- \* The manuscript must be submitted in MS-WORD file format.

#### **INSTRUCTIONS TO AUTHORS**

#### Submission

Submit manuscripts as e-mail attachment to the Editorial Office at:

textroadjournals@gmail.com or info@textroad.com along with covering letter. A manuscript number will be mailed to the corresponding author same day or within 48 hours. The authors may also suggest two to four reviewers for the manuscript (JBASR may designate other reviewers). There is no page limit. The submitting author takes responsibility for the paper during submission and peer review.

#### **Terms of Submission**

Papers must be submitted on the understanding that they have not been published elsewhere (except in the form of an abstract or as part of a published lecture, review, or thesis) and are not currently under consideration by another journal. The submitting author is responsible for ensuring that the article's publication has been approved by all the other coauthors. All enquiries concerning the publication of accepted papers should be addressed to <u>editor@textroad.com</u>.

#### **Review Process**

All manuscripts are reviewed by an editor and members of the Editorial Board or qualified outside reviewers. Decisions will be made as rapidly as possible, and the journal strives to return reviewers' comments to authors within one or two weeks. The editorial board will re-review manuscripts that are accepted pending revision. It is the goal of the JBASR to publish manuscripts within 4 weeks after submission.

#### Style of Manuscripts

Manuscripts should be written in clear, concise and grammatically correct English (with 10 font size and Times New Roman font style) so that they are intelligible to the professional reader who is not a specialist in any particular field. Manuscripts that do not conform to these requirements and the following manuscript format may be returned to the author prior to review for correction. The entire manuscript, including references, should be typed single spaced on one side of the paper. All pages should be numbered consecutively in the bottom centre starting from the title page. The manuscript should be presented in the following order.

#### **Title and Authorship Information**

The title should be a brief phrase (capitalize first letter of each word in the title) describing the contents of the paper. The Title Page should include the authors' full names and affiliations, the name of the corresponding author along with phone, fax and E-mail information. Present addresses of authors should appear as a footnote.

#### Abstract

All manuscripts should not exceed 250-300 words and should describe the scope, hypothesis or rationale for the work and the main findings. Complete sentences, active verbs, and the abstract should be written in the past tense. Standard nomenclature should be used and abbreviations should be avoided. No literature should be cited.

#### Keywords

Key words (5-7 words) should be provided below the Abstract to assist with indexing of the article. These should not duplicate key words from the title.

#### Introduction

This section should include sufficient background information, provide a clear statement of the problem, the relevant literature on the subject, and the proposed approach or solution. The aims of the manuscript should be clearly stated. The introduction should not contain either findings or conclusions. It should be understandable to colleagues from a broad range of scientific disciplines.

#### **Materials and Methods**

This should be complete enough to provide sufficient detail to allow the work to be repeated by others. However, only truly new procedures should be described in detail; previously published procedures should be cited, and important modifications of published procedures should be mentioned briefly. Capitalize trade names and include the manufacturer's name and address. Subheadings should be used. Methods in general use need not be described in detail.

#### Results

Results should be presented in a logical sequence in the text, tables and figures; repetitive presentation of the same data in different forms should be avoided. The results should not contain material appropriate to the Discussion. It should be written in the past tense when describing findings in the authors' experiments. Results should be explained, but largely without referring to the literature.

#### Discussion

The discussion should consider the results in relation to any hypotheses advanced in the Introduction and place the study in the context of other work. Results and Discussion sections can be combined.

#### Conclusions

If an optional conclusion section is used, its content should not substantially duplicate the abstract.

#### Acknowledgment

The acknowledgments of people, grants, funds, etc should be brief.

#### References

Bibliographic references in the text appear like [1, 2, 5, 6], using square brace in superscript. References should be numbered consecutively, with style:

#### Journal paper:

1. Hadjibabaie, M., N. Rastkari, A.Rezaie and M. Abdollahi, 2005. The Adverse Drug Reaction in the Gastrointestinal Tract: An Overview. Intl. J. Pharmacol., 1 (1): 1-8.

#### **Books:**

1. Daniel A. Potter, 2002. Destructive turfgrass insects: Biology, diagnosis and control. Wiley Canada Publishers, pp: 24-67.

#### **Chapters in Book:**

1. Bray R.A., 1994. The leucaena psyllid. In: Forage Tree Legumes in Tropical Agriculture (eds R.C. Gutteridge and H.M. Shelton) pp. 283–291. CAB International, Oxford.

Titles of journals should be given in full. 'In press' can only be used to cite manuscripts actually accepted for publication in a journal. Citations such as 'manuscript in preparation' or 'manuscript submitted' are not permitted. Data from such manuscripts can only be mentioned in the text as 'unpublished data'.

#### A Report:

1. Makarewicz, J.C., T. Lewis and P. Bertram, 1995. Epilimnetic phytoplankton and zooplankton biomass and species composition in Lake Michigan, 1983-1992. U.S. EPA Great Lakes National Program, Chicago, IL. EPA 905-R-95-009.

#### **Conference Proceedings:**

1. Stock, A., 2004. Signal Transduction in Bacteria. In the Proceedings of the 2004 Markey Scholars Conference, pp: 80-89.

#### A Thesis:

1. Strunk, J.L., 1991. The extraction of mercury from sediment and the geochemical partitioning of mercury in sediments from Lake Superior, M. S. thesis, Michigan State Univ., East Lansing, MI.

#### **Tables and Equations**

Tables and equations should not be submitted in a format exceeding the A4 page size (in portrait form). **All tables should be embedded within the manuscript, and must be captioned and numbered sequentially.** Each table should be on a separate page, numbered consecutively in Arabic numerals and supplied with a heading and a legend. Tables should be self-explanatory without reference to the text.

#### Figures / Illustrations / Photographs

Graphics should be supplied as high resolution (at least 300-600 dp.i.) electronic files. Digital images supplied only as low-resolution print-outs cannot be used. Graphs, diagrams, chromatograms, photos, etc. should be prepared as clear, original positives, suitable for reproduction. All figures should be embedded within the manuscript, and must be captioned and numbered sequentially.

#### Proofs

Proofs will be sent via e-mail as an Acrobat PDF file (e-mail attachment) and should be returned within 3 days of receipt. Page proofs are considered to be the final version of the manuscript. With the exception of typographical or minor clerical errors, no changes will be made in the manuscript at the proof stage.

#### **Check List**

We recommend that you ask a colleague to read over your paper prior to submission to ensure it is of a high standard and conforms to a high level of scientific writing.

Before submission of your manuscript, please check that:

- All references cited in the text are included in the reference section.
- All figures and tables are cited in the text.
- Figures are at least 300 d.p.i.
- The pages are numbered.

## Copyright © 2022, TEXTROAD Publishing Corporation







Copyright © 2022, TEXTROAD. All Rights Reserved. TEXTROAD Publishing Corporation